

VU-04/93
UITP 34/1993
hep-th/9404160

$3nj$ -symbols and D -dimensional quantum gravity

P.Demkin¹

*Institute of Theoretical Physics
Uppsala University
Box 803, S-75108
Uppsala, Sweden
paul@rhea.teorfys.uu.se*

Abstract

The model which generalizes Ponzano and Regge $3D$ and Carfora-Martellini-Marzuoli $4D$ euclidean quantum gravity is considered. The euclidean Einstein-Regge action for a D -simplex is given in the semi-classical limit by a gaussian integral of a suitable $3nj$ -symbol.

¹On leave from Department of Physics, Vilnius University, Saulėtekio al.9, 2054, Vilnius, Lithuania

Some time ago G.Ponzano and T.Regge discovered a connection between the expansion of the $6j$ -symbol and the partition function for $3D$ euclidean quantum gravity [1]. These ideas were developed by M.Carfora, M.Martellini and A.Marzuoli for the $4D$ euclidean quantum gravity [2]. In this paper the general correlation between $3nj$ -symbols and D euclidean quantum gravity is shown.

The main aspect here is the correlation between geometrical properties of the D -simplex and the corresponding $3nj$ -symbol. The D -simplex in the D -dimensional space has $D + 1$ vertices and $C_{D+1}^2 = \frac{1}{2}D(D + 1)$ edges. No less than $D - 3$ edges belong to $D - 3$ -dimensional subspace and no more than $\frac{1}{2}D(D - 1) + 3$ edges belong to 3-dimensional subspace R^3 . To every edge belonging to R^3 we relate some quantity of the angular momenta j_i in the $3nj$ -symbol. Therefore, $3nj$ -symbol must contain no less than $\frac{1}{2}D(D - 1) + 3$ angular momenta.

There are $3nj$ -symbols of the first and second kind, depending on the possibility to bound the first and the last momenta [3](see fig.1). We shall consider $3nj$ -symbols of only the first kind. The application of $3nj$ -symbols of the second kind is the same.

Any $3nj$ -symbol can be represented as a sum of products of the corresponding $6j$ -symbols.

$$\left\{ \begin{array}{cccc} j_1 & j_2 & \dots & j_n \\ l_1 & l_2 & \dots & l_n \\ k_1 & k_2 & \dots & k_n \end{array} \right\} = \sum_x (-1)^{R_n + (n-1)x} (2x + 1) \left\{ \begin{array}{ccc} j_1 & k_1 & x \\ k_2 & j_2 & l_1 \end{array} \right\} \times \\ \times \left\{ \begin{array}{ccc} j_2 & k_2 & x \\ k_3 & j_3 & l_2 \end{array} \right\} \times \dots \times \left\{ \begin{array}{ccc} j_{n-1} & k_{n-1} & x \\ k_n & j_n & l_{n-1} \end{array} \right\} \left\{ \begin{array}{ccc} j_n & k_n & x \\ j_1 & k_1 & l_n \end{array} \right\}, \quad (1)$$

where $R_n = \sum_{i=1}^n (j_i + l_i + k_i)$.

Inasmuch as the asymptotic form of the Racah-Wigner $6j$ -symbol according to Ponzano-Regge formula [1] can be expressed by the euclidean Einstein-Regge action $S_R[T]$ for the 3-simplex T (tetrahedron)

$$\left\{ \begin{array}{ccc} j_1 & j_2 & j_3 \\ j_4 & j_5 & j_6 \end{array} \right\} \rightarrow \exp(i \sum_{m=1}^6 j_m \theta_m) = \exp(i S_R[T]), \quad (2)$$

at $j_m \gg 1$, expanding the $3nj$ -symbol into the sum of products of 6j-symbols we would obtain a correlation between the euclidean action of gravity S_R and the $3nj$ -symbol.

There are, however, some difficulties in the implementation of this project. In particular, since expression (2) is valid only at $j_i \gg 1$ expanding the $3nj$ -symbol into 6j-symbols we are summing by the variable moment x , which obeys the condition of triangle $|j_i - j_j| \leq x \leq j_i + j_j$, from which it follows that even at $j_i, j_j \gg 1$, we may receive $|j_i - j_j| \simeq 1$, and expression (2) becomes invalid. Consequently, for any meaning of the space-time dimension D we must choose a special type of the $3nj$ symbol, at the condition

$$3n > \frac{1}{2}D(D-1) + 3, \quad n > 2, \quad D > 3. \quad (3)$$

To escape summing when expanding (1), let us equal one of the momenta to zero:

$$\begin{aligned} & \left\{ \begin{array}{cccc} j_1 & j_2 & \dots & 0 \\ l_1 & l_2 & \dots & l_n \\ k_1 & k_2 & \dots & k_n \end{array} \right\} = \delta(l_n, k_1) \delta(j_{n-1}, l_{n-1}) \times \\ & \times [(2k_1 + 1)(2j_{n-1} + 1)]^{-\frac{1}{2}} (-1)^{R_{n-1} + nk_n - j_1 + k_{n-1} + j_{n-1}} \times \\ & \left\{ \begin{array}{ccc} j_1 & k_1 & x \\ k_2 & j_2 & l_1 \end{array} \right\} \left\{ \begin{array}{ccc} j_2 & k_2 & x \\ k_3 & j_3 & l_2 \end{array} \right\} \dots \left\{ \begin{array}{ccc} j_{n-1} & k_{n-1} & x \\ k_n & j_n & l_{n-1} \end{array} \right\} \end{aligned} \quad (4)$$

In this case the $3nj$ -symbol depends on only $3(n-1)$ momenta.

When the number of edges of the D -simplex belonging to subspace R^3 is a number multiple to 3, the quantity $3n$ in the $3nj$ -symbol is just equal to the number of edges in R^3 plus 3, i.e. $3n = \frac{1}{2}D(D-1) + 6$. When the number of edges in R^3 is the number not multiple to 3, then we put $3n = \frac{1}{2}D(D-1) + 8$ and impose additional conditions like $j_i = j_k$ or/and $j_i = j_{n-1}$ to decrease

the number of independent momenta. Some information about D-simplices and corresponding $3nj$ -symbols for $D \leq 11$ is given in Table 1.

Now, for the reduced $3nj$ -symbol we use equation (1) and the expression of Ponzano-Regge (2):

$$\left\{ \begin{array}{c} \text{reduced} \\ 3nj \end{array} \right\} \mapsto \frac{N}{(12\pi)^{\frac{n-2}{2}} \prod_{k=1}^{\frac{n-2}{2}} V_k^{\frac{1}{2}}} \times \\ \times \exp(i \sum_{m=1}^{n-2} j_m \theta_m + i \sum_{m'=1}^n j_{m'} \theta'_{m'} + i \sum_{m''=2}^n k_{m''} \theta''_{m''}), \quad (5)$$

where $N \equiv N_1 = (-1)^{R_{n-1} + nk_n - j_1 + k_{n-1} + j_{n-1}} \delta(l_n, k_1) \delta(j_{n-1}, l_{n-1}) \times$
 $\times [(2k_1 + 1)(2j_{n-1} + 1)]^{-\frac{1}{2}}$, for $D \neq 3n + 2$ and $N \equiv N_1 \delta(j_i, j_{n-1}) \delta(j_{i-1}, j_{n-2})$,
 $i \neq n$ for $D = 3n + 2$, V_k - volume of the k -th tetrahedron.

Let us introduce new denotations

$$(j_1, \dots, j_{n-1}, l_1, \dots, l_n, k_2, \dots, k_n) \rightarrow (J_1, \dots, J_M), \quad (6)$$

$$(\theta_m, \theta_{m'}, \theta_{m''}) \rightarrow (\psi_1, \dots, \psi_M), \quad (7)$$

Then the reduced $3nj$ -symbol is

$$\left\{ \begin{array}{c} \text{reduced} \\ 3nj \end{array} \right\} \mapsto \frac{N}{(12\pi)^{\frac{n-2}{2}} \prod_{k=1}^{\frac{n-2}{2}} V_k^{\frac{1}{2}}} \times \\ \times \exp(i \sum_{m=1}^M J_m \psi_m). \quad (8)$$

Let us consider the gaussian integral over the real variables ψ_i for the real non singular and symmetric $M \times M$ matrix Δ :

$$Z[J_i, \Delta_{ik}] = \int \left\{ \begin{array}{c} \text{reduced} \\ 3nj \end{array} \right\} \exp(-\frac{1}{2} \sum_{i,k} \psi_i \Delta_{ik} \psi_k) \prod_{j=1}^M d\psi_j \quad . \quad (9)$$

Using the formula for gaussian integration in the semiclassical limit $J_i \gg 1$ we find the following results:

$$Z[J_i, \Delta_{ik}] \sim \frac{\hat{N}}{(\det \Delta)^{\frac{1}{2}}} \exp(-\sum_{i,k=1}^M J_i (\Delta^{-1})_{ik} J_k), \quad (10)$$

where $\hat{N} = (2\pi)^{M-\frac{n-2}{2}} N(6^{-\frac{n-2}{2}} \prod_{k=1}^{\frac{n-2}{2}} V_k^{\frac{1}{2}})^{-1}$.

For the correlation with euclidian quantum gravity we must find the geometrical interpretation of this gaussian integral. The geometrical meaning of expression (9) is more clear if we note that the graphic representation of the $3nj$ -symbol in Fig.1 is a two-dimensional projection of the $2n$ -simplex. Besides, there is a correlation between D and $D+1$ simplices. Every $(D+1)$ -simplex may be built joining two D -simplices, by identification of two $(D-1)$ -subsimplices and linking the two last free vertices by a new edge in $(D+1)$ subspace. This means, in particular, that $(D+1)$ -simplex at $D > 2$ may be represented as a unification of the tetrahedrons with rigid identification edges. This is a geometrical analogue of expanded (7).

Consider

$$\sum_{i,k=1}^M J_i(\Delta^{-1})_{ik} J_k \equiv \sum_{i,k=1}^M \Theta_{ik} A_{ki}, \quad (11)$$

where A_{ki} is the area of the 2-face of D -simplex containing J_i and J_k , and Θ_{ik} is a defect angle if the 2-face containing J_i and J_k belongs to interior of D -simplex, or it is an angle between the outer normals of the two (D_1) -simplices sharing the $(J_i J_k)$ -face, if this 2-face belongs to the boundary of the D -simplex.

For a general D -simplicial manifold M^D with up to an arbitrary term depending on the edge length the euclidean Einstein-Regge action

$$S_R[M^D] \sim \sum_{\sigma \in \text{int} M^D} A(\sigma) \epsilon(\sigma) + \sum_{\sigma \in \partial M^D} A(\sigma) \alpha(\sigma), \quad (12)$$

where σ stands for the 2-simplex where the curvature is concentrated and $A(\sigma)$ is the area of σ ; $\epsilon(\sigma)$ is the defect angle associated with the 2-simplex σ , and $\alpha(\sigma)$ must be interpreted as the angle between the outer normals of the two boundary 3-simplices intersecting at σ .

Then, according to (10) and (11), we may write

$$Z[J_i, \Delta_{ik}] \sim \frac{\hat{N}}{(\det \Delta)^{\frac{1}{2}}} \exp(-S_R[\sigma^D]), \quad (13)$$

where $S_r[\sigma^D]$ is the Einstein-Regge action for the D -simplex σ . We may consider this expression as a semiclassical limit of the partition function for D gravity and as a direct generalization of the Ponzano and Regge and Carfora-Martellini-Marzuoli result.

In principle, there is another way to generalize the Ponzano-Regge ideas. We may try to build $3nj$ -symbols for every unitary algebra $su(N)$ and, possible, for their products and use the correlation between the action of quantum gravity and a corresponding $3nj$ -symbol.

We could receive useful results only for some particular cases in $D = 4$ and $D = 6$ by using the isomorphisms $so(4) \simeq su(2) \oplus su(2)$ and $so(6) \simeq su(4)$. Then, for $su(4)$ algebra, for instance, $21j$ -symbol corresponds to $D = 6$ -simplex. In general, there are several variants for building the euclidean quantum action for higher D as a unification of 3 or 4, or ... $D - 1$ simplices and corresponding $3nj$ - momenta. For other dimensions, we must again use $3nj$ -symbols at $n > 2$. On the other hand, on this way we have one interesting possibility to build a pseudoeuclidean quantum gravity action in $D = 3$ by introducing $6j$ -symbols for $su(1, 1)$ group and use isomorphism $su(1, 1) \simeq so(2, 1)$. In any case we may consider all these possibilities as a kind of selfconsistency: various ways of building the D -simplex must give us the same result.

These ideas may be applied to superspace. By analogy with the rotation case, the super $6j$ -symbol ($s6j$) for the superalgebra $osp(1|2)$ is a symmetric transformation coefficient that relates two bases in an irreducible representation space. The irreducible representations of the superalgebra $osp(1|2)$ have been analysed in [4, 5]. The $s6j$ -symbols for the superalgebra $osp(1|2)$ possess, in addition to the usual tetrahedral symmetry, the so-called Regge symmetry in some particular cases[6]. Then we may introduce $s3nj$ -supersymbols and consider the semiclassical limit of the euclidean action in the corresponding superspace for D supergravity.

Acknowledgements

Author wants to express his gratitude to the Swedish Institute for Grant 304/01 GH/MLH, which gave him the possibility to enjoy the kind hospitality of Prof. Antti Niemi, Doc. Staffan Yngve and all members of the Institute of Theoretical Physics, Uppsala University.

References

- [1] G.Ponzano, T.Regge, Semiclassical limit of Racah coefficients, in: Spectroscopic and group theoretical methods in physics, eds. F.Block et al.(North-Holland, Amsterdam, 1968)pp.1-58.
- [2] M.Carfora, M.Martellini and A.Marzuoli, Phys.Lett.B 299 (1993) 229.
- [3] A.P.Jucys, A.A.Bandzaitis. Theory of angular momentum in quantum mechanics. Mokslas, Vilnius, 1977.
- [4] M.Scheunert, W.Nahm and V.Rittenberg, J.Math.Phys., 18 (1977) 155.
- [5] F.A.Berezin and V.N.Tolstoy, Commun. Math. Phys., 8 (1981) 409.
- [6] M.Daumeus, P.Minnaert, M.Morzymas and S.Toshev, Europhys.Lett., 20 (1992) 671.

D	3	4	5	6	7	10	11	D
$\#V$	4	5	6	7	8	11	12	$D + 1$
$\#E$	6	10	15	21	28	55	66	$1/2D(D + 1)$
$\#E \in R^3$	6	9	13	18	24	48	58	$1/2D(D - 1) + 3$
$3n(3nj)$	6	12	18	21	27	51	63	$\frac{1/2D(D-1)+8,}{1/2D(D-1)+6,} \frac{D=3n+2}{D-others}$
$\#add.cond.$	-	2	4	2	2	2	4	$\frac{4,}{2,} \frac{D=3n+2}{D-others}$

Table 1. Some properties of D -simplices and $3nj$ -symbols.

D is a space-time dimension, $\#V$ is the number of vertices, $\#E$ is the number of edges, $\#E \in R^3$ is the number of edges belonging to subspace R^3 , $3n(3nj)$ is the quantity of the $3nj$ -symbol, $\#add.cond.$ is the number of additional conditions like $j_i = j_k$, which we impose on the corresponding $3nj$ -symbol. One general additional condition $j_n = 0$ is valid for all $3nj$ -symbols to escape summation.